

Appendix B. Summary of Bird Surveys

Table B1. Summary of bird monitoring (for species of conservation importance and/or wetland-dependence) within the Survey Area (excluding the WRA)

| Common Name ⁽⁴⁾ | Scientific Name ⁽⁴⁾ | Conservation Status ⁽²⁾ | Nov Mean ⁽³⁾ | Dec Mean ⁽³⁾ | Jan Mean ⁽³⁾ | Feb Mean ⁽³⁾ | Mar Mean ⁽³⁾ | Apr Mean ⁽³⁾ |
|----------------------------|--------------------------------|------------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| Little Grebe | <i>Tachybaptus ruficollis</i> | LC, (1) | 10.0 | 8.6 | 5.6 | 6.3 | 5.0 | 3.6 |
| Great Cormorant | <i>Phalacrocorax carbo</i> | PRC, (1) | 8.4 | 8.0 | 3.4 | 5.3 | 3.5 | |
| Grey Heron | <i>Ardea cinerea</i> | PRC, (1) | 8.8 | 5.4 | 4.6 | 3.0 | 0.8 | 0.2 |
| Great Egret | <i>Egretta alba</i> | PRC, (RC), (1) | 4.0 | 4.0 | 3.8 | 3.0 | 16.3 | 3.0 |
| Little Egret | <i>Egretta garzetta</i> | PRC, (RC), (1) | 17.8 | 9.8 | 9.6 | 13.5 | 67.5 | 23.2 |
| Eastern Cattle Egret | <i>Bubulcus coromandus</i> | LC, (1) | | | | | 0.3 | 9.0 |
| Chinese Pond Heron | <i>Ardeola bacchus</i> | PRC, (RC), (1) | 9.6 | 4.0 | 8.4 | 6.8 | 6.0 | 7.0 |
| Black-crowned Night Heron | <i>Nycticorax nycticorax</i> | (LC), (1) | | 0.2 | | | | 0.2 |
| Yellow Bittern | <i>Ixobrychus sinensis</i> | LC, (1) | 0.2 | | | | | |
| Black-faced Spoonbill | <i>Platalea minor</i> | PGC, (1) | 0.6 | | V | 2.0 | 2.5 | |
| Northern Shoveler | <i>Anas clypeata</i> | RC, (1) | | 1.2 | 5.8 | | | |
| Tufted Duck | <i>Aythya fuligula</i> | LC, (1) | | 2.8 | 13.0 | 18.3 | 16.3 | |
| Black Kite | <i>Milvus migrans</i> | RC, (1) | | 0.2 | 0.2 | 0.5 | 0.5 | |
| Common Kestrel | <i>Falco tinnunculus</i> | (-) | | V | | | | |
| Peregrine Falcon | <i>Falco peregrinus</i> | LC | | V | | | | |
| White-breasted Waterhen | <i>Amaurornis phoenicurus</i> | (1) | 1.0 | 1.6 | V | 0.5 | 0.3 | 0.6 |
| Common Moorhen | <i>Gallinula chloropus</i> | (1) | 0.8 | 0.4 | | 0.5 | 0.3 | 0.6 |
| Black-winged Stilt | <i>Himantopus himantopus</i> | (RC), (1) | 0.6 | | | | | |
| Pied Avocet | <i>Recurvirostra avosetta</i> | RC, (1) | 4.6 | V | | | | |
| Little Ringed Plover | <i>Charadrius dubius</i> | (LC), (1) | 7.0 | 0.8 | | 1.0 | 1.3 | 0.2 |
| Kentish Plover | <i>Charadrius alexandrinus</i> | RC, (1) | 1.8 | V | | | | |

Proposed Comprehensive Development at Wo Shang Wai Yuen Long

Biannual EM&A Report on Ecology for Nov 2013 - Apr 2014 (Rev A)



| Common Name ⁽⁴⁾ | Scientific Name ⁽⁴⁾ | Conservation Status ⁽²⁾ | Nov Mean ⁽³⁾ | Dec Mean ⁽³⁾ | Jan Mean ⁽³⁾ | Feb Mean ⁽³⁾ | Mar Mean ⁽³⁾ | Apr Mean ⁽³⁾ |
|--------------------------------|---------------------------------|------------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| Green Sandpiper | <i>Tringa ochropus</i> | (1) | | | 0.2 | | | 0.8 |
| Wood Sandpiper | <i>Tringa glareola</i> | (LC), (1) | | | | | | 0.2 |
| Common Sandpiper | <i>Actitis hypoleucos</i> | (1) | 4.0 | 2.0 | 1.6 | 3.5 | 3.0 | 2.6 |
| Common Snipe | <i>Gallinago gallinago</i> | (1) | | | | | | 0.2 |
| Black-headed Gull | <i>Larus ridibundus</i> | PRC, LC, (1) | | | | | 1.3 | |
| Whiskered Tern | <i>Chlidonias hybrida</i> | LC, (1) | | | | 0.8 | 1.5 | 0.2 |
| Eurasian Eagle Owl | <i>Bubo bubo</i> | RC | | | | V | | |
| Pied Kingfisher | <i>Ceryle rudis</i> | (LC), (1) | 0.2 | 0.4 | | 0.3 | 0.8 | 0.4 |
| White-throated Kingfisher | <i>Halcyon smyrnensis</i> | LC, (1) | 0.2 | 0.6 | 0.2 | 0.5 | V | |
| Common Kingfisher | <i>Alcedo atthis</i> | (1) | 2.2 | 1.6 | 1.0 | 0.5 | 1.0 | 2.4 |
| Eastern Yellow Wagtail | <i>Motacilla tschutschensis</i> | (1) | 1.6 | 2.6 | 3.2 | 2.0 | 2.3 | 5.8 |
| White Wagtail | <i>Motacilla alba</i> | (1) | 10.4 | 11.0 | 8.2 | 11.8 | 6.5 | 7.4 |
| Richard's Pipit | <i>Anthus richardi</i> | (1) | | | 1.6 | 0.8 | 1.0 | 1.6 |
| Red-billed Starling | <i>Spodiopsar sericeus</i> | (1) | | 2.6 | 5.4 | 6.8 | 1.3 | 0.8 |
| White-cheeked Starling | <i>Spodiopsar cineraceus</i> | PRC, (1) | 0.2 | | | | | |
| Collared Crow | <i>Corvus torquatus</i> | LC, (1) | 0.8 | 0.4 | | | 0.5 | 1.6 |
| No. of species recorded | | | 22 | 25 | 19 | 22 | 24 | 22 |

Note:

- (1) Indicates wetland-dependent or wetland-associated species.
- (2) Conservation status follows that of Fellowes et al. (2002) and BirdLife International listing (2010). See Table B3.
- (3) Refers to the mean number of individuals recorded in each survey in the Survey Area (excluding the WRA).
- (4) Follows HK bird list (dated 2013-3-25).
- (5) "V" indicates the species is recorded outside regular surveys.

Table B2. Summary of bird monitoring (for species of conservation importance and/or wetland-dependence) in the WRA

| Common Name ⁽⁴⁾ | Scientific Name ⁽⁴⁾ | Conservation Status ⁽²⁾ | Nov Mean ⁽³⁾ | Dec Mean ⁽³⁾ | Jan Mean ⁽³⁾ | Feb Mean ⁽³⁾ | Mar Mean ⁽³⁾ | Apr Mean ⁽³⁾ |
|----------------------------|--------------------------------|------------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| Little Grebe | <i>Tachybaptus ruficollis</i> | LC, (1) | 0.6 | 2.2 | 1.8 | 5.3 | 4.5 | 2.4 |
| Great Cormorant | <i>Phalacrocorax carbo</i> | PRC, (1) | V | 0.4 | 0.6 | 0.5 | | |
| Grey Heron | <i>Ardea cinerea</i> | PRC, (1) | 2.8 | 2.6 | 1.6 | 2.5 | | |
| Great Egret | <i>Egretta alba</i> | PRC, (RC), (1) | 0.6 | 2.4 | 1.2 | 1.8 | 0.8 | 0.2 |
| Intermediate Egret | <i>Egretta intermedia</i> | RC, (1) | | | | V | | |
| Little Egret | <i>Egretta garzetta</i> | PRC, (RC), (1) | 1.4 | 2.4 | 4.4 | 4.0 | 3.8 | 3.2 |
| Eastern Cattle Egret | <i>Bubulcus coromandus</i> | LC, (1) | | | | | | 1.6 |
| Chinese Pond Heron | <i>Ardeola bacchus</i> | PRC, (RC), (1) | 1.4 | 2.6 | 6.8 | 3.5 | 2.5 | 1.0 |
| Black-crowned Night Heron | <i>Nycticorax nycticorax</i> | (LC), (1) | | | | | V | |
| Yellow Bittern | <i>Ixobrychus sinensis</i> | LC, (1) | 0.4 | 0.2 | | | V | V |
| Black-faced Spoonbill | <i>Platalea minor</i> | PGC, (1) | | | | V | V | |
| Northern Shoveler | <i>Anas clypeata</i> | RC, (1) | | 1.2 | 7.0 | 5.5 | V | |
| Black Kite | <i>Milvus migrans</i> | RC, (1) | 0.2 | | V | 0.5 | 0.8 | 0.4 |
| Crested Serpent Eagle | <i>Spilornis cheela</i> | (LC) | | | | V | | |
| White-breasted Waterhen | <i>Amaurornis phoenicurus</i> | (1) | 0.4 | | | | V | 0.2 |
| Common Moorhen | <i>Gallinula chloropus</i> | (1) | | | | | V | 0.2 |
| Black-winged Stilt | <i>Himantopus himantopus</i> | (RC), (1) | V | | | | | |
| Little Ringed Plover | <i>Charadrius dubius</i> | (LC), (1) | 0.4 | | V | 0.5 | 0.8 | |
| Common Greenshank | <i>Tringa nebularia</i> | LC, (1) | | V | | | V | |
| Green Sandpiper | <i>Tringa ochropus</i> | (1) | 0.6 | V | V | | 0.5 | |
| Wood Sandpiper | <i>Tringa glareola</i> | (LC), (1) | V | V | V | V | V | |
| Common Sandpiper | <i>Actitis hypoleucos</i> | (1) | 1.6 | 0.2 | V | 0.3 | 1.3 | 0.2 |
| Pintail Snipe | <i>Gallinago stenura</i> | (1) | V | | | | | |
| Eurasian Eagle Owl | <i>Bubo bubo</i> | RC | V | | | V | | |
| Pied Kingfisher | <i>Ceryle rudis</i> | (LC), (1) | | 0.4 | V | 0.3 | V | |

Proposed Comprehensive Development at Wo Shang Wai Yuen Long

Biannual EM&A Report on Ecology for Nov 2013 - Apr 2014 (Rev A)



| Common Name ⁽⁴⁾ | Scientific Name ⁽⁴⁾ | Conservation Status ⁽²⁾ | Nov Mean ⁽³⁾ | Dec Mean ⁽³⁾ | Jan Mean ⁽³⁾ | Feb Mean ⁽³⁾ | Mar Mean ⁽³⁾ | Apr Mean ⁽³⁾ |
|--------------------------------|---------------------------------|------------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| White-throated Kingfisher | <i>Halcyon smyrnensis</i> | LC, (1) | | 0.2 | | | | |
| Common Kingfisher | <i>Alcedo atthis</i> | (1) | 2.0 | 1.6 | 1.2 | 2.3 | 0.8 | 1.0 |
| Eastern Yellow Wagtail | <i>Motacilla tschutschensis</i> | (1) | 0.2 | 0.6 | 0.6 | 1.0 | 0.3 | 0.2 |
| White Wagtail | <i>Motacilla alba</i> | (1) | 3.6 | 2.6 | 1.2 | 1.3 | 2.5 | 1.6 |
| Richard's Pipit | <i>Anthus richardi</i> | (1) | | 0.4 | 0.8 | 2.0 | 2.3 | 3.0 |
| Zitting Cisticola | <i>Cisticola juncidis</i> | LC, (1) | | | 0.4 | | 0.3 | 1.2 |
| Chinese Penduline Tit | <i>Remiz consobrinus</i> | RC, (1) | | V | | 4.5 | V | |
| Red-billed Starling | <i>Spodiopsar sericeus</i> | (1) | | | | V | | |
| Collared Crow | <i>Corvus torquatus</i> | LC, (1) | 0.2 | | | V | 0.5 | 0.8 |
| No. of species recorded | | | 21 | 20 | 18 | 22 | 24 | 16 |

Note:

(1) Indicates wetland-dependent or wetland-associated species.

(2) Conservation status follows that of Fellowes et al. (2002) and BirdLife International listing (2010). See Table B3.

(3) Refers to the mean number of individuals recorded in each survey in the Survey Area (excluding the WRA)

(4) Follows HK bird list (dated 2013-3-25)

V Indicates the species is recorded outside regular surveys.

Table B3. Conservation Status Categories of Bird Species

| Code | Category | Brief/Description | Source |
|------|----------------------------|---|-------------------------------|
| GC | Global Concern | Habitat loss/damage in Hong Kong would pose significant threat to global survival | Fellowes <i>et al.</i> (2002) |
| RC | Regional Concern | Habitat loss/damage in Hong Kong would pose significant threat to regional survival. | |
| LC | Local Concern | Habitat loss/damage in Hong Kong would pose significant threat to local survival. | |
| PGC | Potential Global Concern | Large, secure population in Hong Kong is of global significance. | |
| PRC | Potential Regional Concern | Large, secure population in Hong Kong is of regional significance. | |
| CR | Critically Endangered | Best available evidence indicates that it meets any of the criteria A to E for Critically Endangered, and it is therefore considered to be facing an extremely high risk of extinction in the wild. | BirdLife International (2010) |
| EN | Endangered | Best available evidence indicates that it meets any of the criteria A to E for Endangered, and it is therefore considered to be facing a very high risk of extinction in the wild. | |
| VU | Vulnerable | Best available evidence indicates that it meets any of the criteria A to E for Vulnerable, and it is therefore considered to be facing a high risk of extinction in the wild. | |
| NT | Near Threatened | Does not qualify for Critically Endangered, Endangered or Vulnerable now, but is close to qualifying for or is likely to qualify for a threatened category in the near future. | |

Table B4. Summary of Bird Species Recorded at the WRA and their Respective Monthly Mean between September 2010 and October 2012

| Common name | Scientific name | Conservation status | Sep-10 | Oct-10 | Nov-10 | Dec-10 | Jan-11 | Feb-11 | Mar-11 | Apr-11 | May-11 | Jun-11 | Jul-11 | Aug-11 | Sep-11 | Oct-11 | Nov-11 | Dec-11 | Jan-12 | Feb-12 | Mar-12 | Apr-12 | May-12 | Jun-12 | Jul-12 | Aug-12 | Sep-12 | Oct-12 |
|---------------------------|---------------------------------|---------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Little Grebe | <i>Tachybaptus ruficollis</i> | LC, (1) | | | | | 0.5 | | | | | | 1.0 | 3.8 | 0.8 | 0.5 | 0.5 | | 0.6 | | 1.4 | 1.6 | 0.8 | 1.0 | 1.3 | 0.2 | | 0.2 |
| Great Cormorant | <i>Phalacrocorax carbo</i> | PRC, (1) | | | | | 0.8 | | | | | | | | | 0.3 | 0.3 | 0.4 | 1.2 | 0.7 | 0.2 | | | | | | | |
| Grey Heron | <i>Ardea cinerea</i> | PRC, (1) | | | 1.0 | 1.6 | 2.0 | 2.0 | 0.4 | | | | | | | 1.0 | 2.5 | 1.8 | 0.8 | 2.3 | 0.8 | | | | 0.3 | | | 0.4 |
| Purple Heron | <i>Ardea purpurea</i> | RC, (1) | | | | | | | | | | | | | | 0.3 | 0.3 | | | | | | | | | | | |
| Great Egret | <i>Ardea alba</i> | PRC, (1) | | | | 0.4 | 14.0 | 2.0 | 0.2 | 0.5 | 0.3 | 0.5 | 0.3 | | 1.5 | 1.8 | 0.8 | 1.2 | 0.6 | 0.7 | 0.6 | 0.6 | 1.0 | 0.3 | | 0.4 | 0.5 | 1.5 |
| Intermediate Egret | <i>Egretta intermedia</i> | RC, (1) | | | | | 1.7 | 0.8 | 0.2 | | | | | | 1.5 | 1.5 | 1.5 | 1.0 | 0.6 | 0.7 | 0.6 | 0.6 | | | | | 0.3 | 0.9 |
| Little Egret | <i>Egretta garzetta</i> | PRC, (1) | | | | 1.8 | 8.0 | 2.0 | 0.4 | 0.5 | 0.8 | 0.3 | 0.3 | 1.0 | 1.0 | 0.5 | 1.5 | 1.6 | 1.4 | 2.0 | 1.6 | | | | | 0.8 | 0.3 | 1.2 |
| Eastern Cattle Egret | <i>Bubulcus coromandus</i> | (LC), (1) | | | | | | | | | | 0.3 | 11.8 | | 1.3 | 0.3 | | | 0.4 | | | | | 0.3 | | 0.2 | | |
| Chinese Pond Heron | <i>Ardeola bacchus</i> | PRC, (1) | | | 1.0 | 0.2 | | | | 0.3 | 0.3 | 0.3 | 1.5 | 4.2 | 7.8 | 4.5 | 5.3 | 2.4 | 0.3 | 2.3 | 2.6 | 1.4 | 0.8 | 1.0 | 0.8 | 2.0 | 1.8 | 2.2 |
| Yellow Bittern | <i>Ixobrychus sinensis</i> | (LC), (1) | | | | | | | | | | | | | | 0.3 | 0.3 | | | | | | | | | | | |
| Black-crowned Night Heron | <i>Nycticorax nycticorax</i> | LC, (1) | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Eurasian Spoonbill | <i>Platalea leucorodia</i> | LC, (1) | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Black-faced Spoonbill | <i>Platalea minor</i> | PGC, EN, (1) | | | | | 0.3 | 10.3 | | | | | | | | | | | 0.4 | | | | | | | | | |
| Mandarin Duck | <i>Aix galericulata</i> | (1) | | | | | | | | | | | | | | 0.3 | | | | | | | | | | | | |
| Eurasian Wigeon | <i>Anas penelope</i> | RC, (1) | | | | 1.6 | 2.5 | | | | | | | | | | | | | | | | | | | | | |
| Eurasian Teal | <i>Anas crecca</i> | RC, (1) | | | | | 1.0 | | | | | | | | | | | | | | | | | | | | | |
| Northern Pintail | <i>Anas acuta</i> | RC, (1) | | | | 0.2 | 7.0 | | | | | | | | | | | | | | | | | | | | | |
| Northern Shoveler | <i>Anas clypeata</i> | RC, (1) | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Tufted Duck | <i>Aythya fuligula</i> | LC, (1) | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Garganey | <i>Anas querquedula</i> | (1) | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Western Osprey | <i>Pandion haliaetus</i> | RC, (1) | | | | 0.2 | | | | | | | | | | | | | | | | | | | | | | |
| Black Kite | <i>Milvus migrans</i> | (RC) | | | | | 0.8 | 0.3 | 0.2 | | 0.3 | | | | 0.3 | | | | 0.2 | | | 0.2 | 0.5 | | | | | 0.4 |
| Crested Serpent Eagle | <i>Spilornis cheela</i> | (LC) | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Eastern Buzzard | <i>Buteo japonicus</i> | - | | | | | | | 0.2 | | | | | | | | | | | | | | | | | | | |
| Common Kestrel | <i>Falco tinnunculus</i> | - | | | 1.0 | 0.4 | 1.0 | 0.3 | 0.4 | | | | | | | | | | | | | | | | | | | |
| Eurasian Hobby | <i>Falco subbuteo</i> | (LC), (1) | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Peregrine Falcon | <i>Falco peregrinus</i> | - | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Imperial Eagle | <i>Aquila heliaca</i> | - | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Japanese Quail | <i>Coturnix japonica</i> | LC, (1) | | | | | | | | | | | | | | | | | | | | | | | | | | |
| White-breasted Waterhen | <i>Amauromis phoenicurus</i> | (1) | | | | | | | | | 0.5 | 0.8 | 1.3 | 2.6 | 1.3 | 1.5 | 0.3 | | | 0.4 | | | | 0.5 | 1.0 | 0.5 | | |
| Common Moorhen | <i>Gallinula chloropus</i> | (1) | | | | | | | | | | | | | | | | | | | | | | | 0.2 | | | |
| Pheasant-tailed Jacana | <i>Hydrophasianus chirurgus</i> | LC, (1) | | | | | | | | | | | | | | | 0.3 | | | | | | | | | | | |
| Greater Painted-snipe | <i>Rostratula benghalensis</i> | LC, (1) | | | | | | | | 0.5 | | | | | | | | | | | | | | | | | | |
| Black-winged Stilt | <i>Himantopus himantopus</i> | RC, (1) | 8.0 | | | | | | | | | | | | | | | | 1.0 | | | | | | | | | |
| Pied Avocet | <i>Recurvirostra avosetta</i> | RC, (1) | | | | 0.2 | | | | | | | | | | | | | | | 0.2 | | | | | | | |
| Oriental Pratincole | <i>Glareola maldivarum</i> | LC, (1) | | | | | | | | 2.5 | | | | | | | | | | | | | | | | | | |
| Little Ringed Plover | <i>Charadrius dubius</i> | LC, (1) | 3.0 | | 24.5 | 7.4 | 3.8 | 9.0 | 6.8 | 10.5 | 5.5 | 0.5 | | 0.2 | | | | | | 1.0 | 0.6 | | | | | | | |

| Common name | Scientific name | Conservation status | Sep-10 | Oct-10 | Nov-10 | Dec-10 | Jan-11 | Feb-11 | Mar-11 | Apr-11 | May-11 | Jun-11 | Jul-11 | Aug-11 | Sep-11 | Oct-11 | Nov-11 | Dec-11 | Jan-12 | Feb-12 | Mar-12 | Apr-12 | May-12 | Jun-12 | Jul-12 | Aug-12 | Sep-12 | Oct-12 |
|---------------------------|------------------------------------|---------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Kentish Plover | <i>Charadrius alexandrinus</i> | RC, (1) | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Spotted Redshank | <i>Tringa erythropus</i> | RC, (1) | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Common Redshank | <i>Tringa totanus</i> | RC, (1) | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Marsh Sandpiper | <i>Tringa stagnatilis</i> | RC, (1) | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Common Greenshank | <i>Tringa nebularia</i> | RC, (1) | | | 0.5 | | | | | | | | | | | | 0.3 | 0.8 | 0.2 | | | | | | | | | |
| Green Sandpiper | <i>Tringa ochropus</i> | (1) | | | 1.0 | 0.4 | 0.5 | 0.5 | 0.4 | | | | 0.3 | | | | | | 0.2 | | 0.2 | 0.4 | | | | 0.2 | | |
| Wood Sandpiper | <i>Tringa glareola</i> | LC, (1) | | | | | | 0.5 | 1.2 | | | | | | | | | | 0.8 | 0.3 | | | | | | 0.6 | | |
| Common Sandpiper | <i>Actitis hypoleucos</i> | (1) | 1.0 | | 0.5 | 1.0 | | 0.5 | 0.6 | 1.0 | 0.3 | | | 0.2 | | | | | | | | 0.4 | 0.2 | | | | | 0.9 |
| Eurasian Woodcock | <i>Scolopax rusticola</i> | (1) | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pintail/Swinhoe's Snipe* | <i>Gallinago stenura/G. megala</i> | LC*,(1) | | | 0.5 | | | | | | | | | | | 0.3 | | | | | | | | | | | | |
| Common Snipe | <i>Gallinago gallinago</i> | (1) | | | | | | 0.3 | 0.2 | | | | | | | | | | | | | | | | | | | |
| Red-necked Stint | <i>Calidris ruficollis</i> | LC, (1) | 3.0 | | | | | | | | | | | | | | | | | | | | | | | | | |
| Temminck's Stint | <i>Calidris temminckii</i> | LC, (1) | | | 0.5 | | | 2.5 | 0.4 | 1.3 | | | | | | | | | | | | | | | | | | |
| Long-toed Stint | <i>Calidris subminuta</i> | LC, (1) | | | | | | | | | | | | | | | | | | 0.3 | | | | | | | | |
| Whiskered Tern | <i>Chlidonias hybrida</i> | (1) | | | | | | | | | | | | | | | | | | | | | 1.3 | | | | | |
| Domestic Pigeon | <i>Columba livia</i> | - | | | | | | | | | | | | DS | | | | | | | | | | | | | | |
| Oriental Turtle Dove | <i>Streptopelia orientalis</i> | - | | | | | | | | | | | | | 0.2 | | | | | | | | | | | | | |
| Red Turtle Dove | <i>Streptopelia tranquebarica</i> | - | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Spotted Dove | <i>Spilopelia chinensis</i> | - | | | | | | | | | | | | | | | | DS | | DS | DS | | DS | | | DS | | 0.6 |
| Asian Koel | <i>Eudynamis scolopaceus</i> | - | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Greater Coucal# | <i>Centropus sinensis</i> | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Eurasian Eagle Owl | <i>Bubo bubo</i> | RC | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Savanna Nightjar | <i>Caprimulgus affinis</i> | - | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pacific Swift | <i>Apus pacificus</i> | (LC) | | | | | | | | 0.3 | | | | | | | | | | | | | | | | | | |
| House Swift | <i>Apus nipalensis</i> | - | 10.0 | | | | | | 0.2 | | | | | | | | | | | DS | DS | | | | | | | |
| Pied Kingfisher | <i>Ceryle rudis</i> | (LC), (1) | | | | 0.2 | 0.3 | 0.5 | 0.2 | | | 0.3 | | | | | | | | | DS | DS | | | 0.5 | | | |
| Common Kingfisher | <i>Alcedo atthis</i> | (1) | | | | 0.4 | 0.5 | 0.3 | | 0.3 | 0.5 | 1.0 | 0.3 | 1.0 | 0.8 | 0.5 | 0.8 | 1.0 | 0.4 | | 0.8 | | | | | 0.6 | | 0.4 |
| White-throated Kingfisher | <i>Halcyon smyrnensis</i> | (LC) | | | | | | | | | | | | | | | 0.3 | | | | | | | | 0.3 | | | |
| Barn Swallow | <i>Hirundo rustica</i> | - | 25.0 | | | | | 1.5 | | | | | | | | | | | | | DS | DS | | DS | | DS | DS | 3.0 |
| Red-rumped Swallow | <i>Hirundo daurica</i> | - | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Eastern Yellow Wagtail | <i>Motacilla tschutschensis</i> | (1) | | | 14.5 | 10.2 | 15.0 | 23.5 | 8.2 | 11.8 | 1.8 | | | 0.2 | 0.8 | 1.5 | 2.0 | 1.2 | 0.6 | | 4.0 | 1.0 | | | | | | |
| Grey Wagtail | <i>Motacilla cinerea</i> | (1) | | | | 0.2 | | 0.3 | 0.2 | | | | | | 0.3 | | | | | | | | | | | | | |
| White Wagtail | <i>Motacilla alba</i> | (1) | 3.0 | | | 1.8 | | | | | | | | | | | | DS | DS | DS | DS | DS | | | | DS | 1.8 | 3.2 |
| Richard's Pipit | <i>Anthus richardi</i> | (1) | | | 1.5 | 0.8 | 0.5 | 1.0 | 1.0 | 1.8 | 0.5 | | | | | 0.3 | | DS | | DS | DS | DS | | | | | | |
| Olive-backed Pipit | <i>Anthus hodgsoni</i> | - | | | 0.5 | | | | | | | | | | | | | DS | | | | | | | | | | |
| Red-throated Pipit | <i>Anthus cervinus</i> | LC | | 2.0 | 2.5 | 1.2 | 0.5 | 0.5 | 0.2 | | | | | | | | | | | | | 0.4 | | | | | | 0.7 |
| Buff-bellied Pipit | <i>Anthus rubescens</i> | LC | | 1.0 | 1.5 | | | | | | | | | | | | | | | | | | | | | | | |
| Red-whiskered Bulbul | <i>Pycnonotus jocosus</i> | - | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Chinese Bulbul | <i>Pycnonotus sinensis</i> | - | | | | | | | | | | | | | | | | | | | DS | | DS | | | DS | | |
| Long-tailed Shrike | <i>Lanius schach</i> | - | | | | | | | | | | | | | | | | | | DS | DS | | | | | DS | DS | |
| Oriental Magpie Robin | <i>Copsychus saularis</i> | - | | | | | | | | | | | | | | | | | | | DS | | | | | | | |
| Daurian Redstart | <i>Phoenicurus aureus</i> | - | | | | | | | | | | | | | | | | | | | | | | | | | | |

| Common name | Scientific name | Conservation status | Sep-10 | Oct-10 | Nov-10 | Dec-10 | Jan-11 | Feb-11 | Mar-11 | Apr-11 | May-11 | Jun-11 | Jul-11 | Aug-11 | Sep-11 | Oct-11 | Nov-11 | Dec-11 | Jan-12 | Feb-12 | Mar-12 | Apr-12 | May-12 | Jun-12 | Jul-12 | Aug-12 | Sep-12 | Oct-12 |
|--------------------------------|----------------------------------|---------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Stejneger's Stonechat | <i>Saxicola stejnegeri</i> | - | | 3.0 | 0.5 | 1.0 | 1.0 | 0.8 | 0.2 | 0.3 | | | | | | 1.0 | | DS | DS | | DS | | | | | | | |
| Masked Laughingthrush | <i>Garrulax perspicillatus</i> | - | | | | | | | | | | | | | | | | | | | | | | | DS | | | |
| Lanceolated Warbler | <i>Locustella lanceolata</i> | (1) | | | | | | | | | | | | | | 0.3 | | | | | | | | | | | | |
| Pallas's Grasshopper Warbler | <i>Locustella certhiola</i> | LC, (1) | | | | | | | | | | | | | | 0.3 | | | | | | | | | | | | |
| Black-browed Reed Warbler | <i>Acrocephalus bistrigiceps</i> | (1) | | | | | | | | | | | | | | 0.3 | | | | | | | | | | | | |
| Zitting Cisticola | <i>Cisticola juncidis</i> | LC | | | | | | 0.2 | 1.5 | 2.5 | 2.5 | 1.0 | 0.4 | | 0.3 | 0.8 | 1.8 | 0.2 | 0.3 | 0.6 | | | | | | 0.2 | | |
| Yellow-bellied Prinia | <i>Prinia flaviventris</i> | - | | | | | | | | | | | | | | | | DS | | DS | | DS | | DS | DS | 0.8 | 0.4 | |
| Plain Prinia | <i>Prinia inornata</i> | - | | | | | | | | | | | | | | | | | | DS | | DS | | DS | DS | | | |
| Dusky Warbler | <i>Phylloscopus fuscatus</i> | - | | | | | | | | | | | | | | 0.5 | | DS | | | | | | | | | | |
| Yellow-browed warbler | <i>Phylloscopus inornatus</i> | - | | | | | | | | | | | | | | | | | | | | | | | | | | 1.4 |
| Chinese Penduline-Tit | <i>Remiz consobrinus</i> | RC, (1) | | | | | | | | | | | | | | | | | | | 1.2 | 0.2 | | | | | | |
| Japanese White-eye | <i>Zosterops japonica</i> | - | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Little Bunting | <i>Emberiza pusilla</i> | - | | | 0.5 | | | | | | | | | | | | | | | | | | | | | | | |
| Black-faced Bunting | <i>Emberiza spodocephala</i> | - | | | | | | | | 0.5 | | | | | | | | | | | | | | | | | | |
| Chinese Grosbeak | <i>Eophona migratoria</i> | LC | | | | | | | | | | | | | | 0.3 | | | | | | | | | | | | |
| Scaly-breasted Munia | <i>Lonchura punctulata</i> | - | 20.0 | | | | 2.5 | | 15.0 | 7.5 | | | | | | | | | | | | DS | | DS | DS | DS | 1.5 | |
| Eurasian Tree Sparrow | <i>Passer montanus</i> | - | | | | | | | | | | | | | | | | | | | DS | | DS | | DS | | | |
| Red-billed Starling | <i>Spodiopsar sericeus</i> | GC | | | | | | | | | | | | | 0.3 | | 0.8 | 1.6 | 6.6 | 39.0 | 17.4 | | | | | | | |
| White-cheeked Starling | <i>Spodiopsar cineraceus</i> | PRC | | | | | | | | | | | | 6.6 | | | | | | | | | | | | | | |
| Black-collared Starling | <i>Gracupica nigricollis</i> | - | | | | | | | | | | | | | | | | DS | DS | DS | DS | | | | | DS | | |
| White-shouldered Starling | <i>Sturnia sinensis</i> | (LC) | | | | | | | | 0.3 | | 0.5 | 2.4 | | | | | | | | | | | | | | | |
| Common Myna | <i>Acridotheres tristis</i> | (1) | | | | | | | | | | | | | | | | | | DS | | | | | | | | |
| Crested Myna | <i>Acridotheres cristatellus</i> | - | 3.0 | | | | | | | | | | | | | | | DS | | | | DS | | | DS | | DS | V |
| Black-napped Oriole | <i>Oriolus chinensis</i> | LC | | | | | | | | | | | | | 1.0 | | | | | | | | | | | | | |
| Black Drongo | <i>Dicrurus macrocercus</i> | - | | | | | | | | | | | | | | | | | | | | | | | DS | DS | | |
| Azure-winged Magpie | <i>Cyanopica cyanus</i> | - | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Eurasian Magpie | <i>Pica pica</i> | - | | | | | | | | | | | | | | | | | | | DS | | | | | DS | 0.3 | |
| Large-billed Crow | <i>Corvus macrorhynchos</i> | - | | | | 0.4 | | | 0.2 | | | | | | | | | | | | | | | | | | | |
| Collared Crow | <i>Corvus torquatus</i> | LC, NT | | | | 0.2 | 0.3 | | | 0.5 | | 0.3 | | 0.8 | 0.8 | 0.8 | 0.5 | | | | | | | 0.8 | | | | |
| No. of species recorded | | 108 | | | | | | | | | | | | | | | | | | | | | | | | | | |

Note: Conservation Status follows that of Fellowes et. al. (2002)

(1) Indicates the bird species is wetland dependent.

* Pintail Snipe and Swinhoe's Snipe cannot be distinguished in field, conservation status refers to Swinhoe's Snipe.

V Indicates the species is recorded outside regular surveys.

DS Indicates species recorded during survey.

Greater Coucal is listed as vulnerable (VU) in China Red Data Book and it is protected under terrestrial wildlife state protection (category II).

Table B5. Summary of Bird Species Recorded at the WRA and their Respective Monthly Mean between November 2012 and April 2014

| Common name | Scientific name | Conservation status | Nov-12 | Dec-12 | Jan-13 | Feb-13 | Mar-13 | Apr-13 | May-13 | Jun-13 | Jul-13 | Aug-13 | Sep-13 | Oct-13 | Nov-13 | Dec-13 | Jan-14 | Feb-14 | Mar-14 | Apr-14 |
|---------------------------|---------------------------------|---------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Little Grebe | <i>Tachybaptus ruficollis</i> | LC, (1) | 1.0 | 1.0 | 1.2 | 0.4 | 0.6 | 0.4 | 1.5 | 0.5 | | 0.6 | 0.5 | 1.2 | 0.6 | 2.2 | 1.8 | 5.3 | 4.5 | 2.4 |
| Great Cormorant | <i>Phalacrocorax carbo</i> | PRC, (1) | 0.8 | 3.2 | 0.8 | 1.0 | 0.0 | 0.0 | | | | | | 0.2 | V | 0.4 | 0.6 | 0.5 | | |
| Grey Heron | <i>Ardea cinerea</i> | PRC, (1) | 0.8 | 1.4 | 0.8 | 1.4 | 0.0 | 0.0 | | | | | V | 1.0 | 2.8 | 2.6 | 1.6 | 2.5 | | |
| Purple Heron | <i>Ardea purpurea</i> | RC, (1) | | | | | | | | | | | | | | | | | | |
| Great Egret | <i>Ardea alba</i> | PRC, (1) | 1.4 | 0.6 | 1.0 | 0.8 | 0.2 | 0.6 | | 0.8 | 0.6 | | | 0.6 | 0.6 | 2.4 | 1.2 | 1.8 | 0.8 | 0.2 |
| Intermediate Egret | <i>Egretta intermedia</i> | RC, (1) | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | | | | | | | | | | V | | |
| Little Egret | <i>Egretta garzetta</i> | PRC, (1) | 0.8 | 2.0 | 2.2 | 1.4 | 0.6 | 1.2 | 0.6 | 1.8 | 0.4 | 3.4 | 0.5 | 2.2 | 1.4 | 2.4 | 4.4 | 4.0 | 3.8 | 3.2 |
| Eastern Cattle Egret | <i>Bubulcus coromandus</i> | (LC), (1) | | | | | | | | | | | | | | | | | | 1.6 |
| Chinese Pond Heron | <i>Ardeola bacchus</i> | PRC, (1) | 1.6 | 2.2 | 2.4 | 0.4 | 0.0 | 0.0 | | | 0.8 | 2.0 | 0.5 | 1.4 | 1.4 | 2.6 | 6.8 | 3.5 | 2.5 | 1.0 |
| Yellow Bittern | <i>Ixobrychus sinensis</i> | (LC), (1) | | | | | | V | 1.0 | 2.5 | 0.2 | 1.4 | 1.3 | 0.8 | 0.4 | 0.2 | | | V | V |
| Black-crowned Night Heron | <i>Nycticorax nycticorax</i> | LC, (1) | V | | | 0.2 | | | | | | | | V | | | | | V | |
| Eurasian Spoonbill | <i>Platalea leucorodia</i> | LC, (1) | | | | | | | | | | | | | | | | | | |
| Black-faced Spoonbill | <i>Platalea minor</i> | PGC, EN, (1) | | | | | | | | | | | | | | | | V | V | |
| Mandarin Duck | <i>Aix galericulata</i> | (1) | | | | | | | | | | | | | | | | | | |
| Eurasian Wigeon | <i>Anas penelope</i> | RC, (1) | | | | | | | | | | | | | | | | | | |
| Eurasian Teal | <i>Anas crecca</i> | RC, (1) | | | | | | | | | | | | | | | | | | |
| Northern Pintail | <i>Anas acuta</i> | RC, (1) | | | | | | | | | | | | | | | | | | |
| Northern Shoveler | <i>Anas clypeata</i> | RC, (1) | | | V | | | | | | | | | | | 1.2 | 7.0 | 5.5 | V | |
| Tufted Duck | <i>Aythya fuligula</i> | LC, (1) | | | V | | | | | | | | | | | | | | | |
| Garganey | <i>Anas querquedula</i> | (1) | | | | | | | | | | | | | | | | | | |
| Western Osprey | <i>Pandion haliaetus</i> | RC, (1) | | | | | | | | | | | | | | | | | | |
| Black Kite | <i>Milvus migrans</i> | (RC) | 0.0 | 0.2 | 0.0 | 0.0 | 0.6 | 0.0 | 0.2 | 0.3 | | 1.2 | 0.8 | 0.2 | 0.2 | | V | 0.5 | 0.8 | 0.4 |
| Crested Serpent Eagle | <i>Spilornis cheela</i> | (LC) | | | | | | | | | | | | | | | | V | | |
| Eastern Buzzard | <i>Buteo japonicus</i> | - | | | | | | | | | | | | | | | | | | |
| Common Kestrel | <i>Falco tinnunculus</i> | - | | | | | | | | | | | | | | | | | | |
| Eurasian Hobby | <i>Falco subbuteo</i> | (LC), (1) | | | | | | | | | | | | | | | | | | |
| Peregrine Falcon | <i>Falco peregrinus</i> | - | | V | | | | | | | | | | | | V | | | | |
| Imperial Eagle | <i>Aquila heliaca</i> | - | | | V | | | | | | | | | | | | | | | |
| Japanese Quail | <i>Coturnix japonica</i> | LC, (1) | | | | | | | | | | | | V | | | | | | |
| White-breasted Waterhen | <i>Amaurornis phoenicurus</i> | (1) | 0.2 | 0.0 | 0.2 | 0.0 | 0.4 | 0.2 | V | 1.8 | 0.2 | 0.2 | V | 0.8 | 0.4 | | | | V | 0.2 |
| Common Moorhen | <i>Gallinula chloropus</i> | (1) | | | | | | | | | | | | | | | | | V | 0.2 |
| Pheasant-tailed Jacana | <i>Hydrophasianus chirurgus</i> | LC, (1) | | | | | | | | | | | | | | | | | | |
| Greater Painted-snipe | <i>Rostratula benghalensis</i> | LC, (1) | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | | | | | | | | | | | | |
| Black-winged Stilt | <i>Himantopus himantopus</i> | RC, (1) | | | | | | | | | 0.2 | 0.6 | V | 0.2 | V | | | | | |
| Pied Avocet | <i>Recurvirostra avosetta</i> | RC, (1) | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | | | | | | | | | | | | |
| Oriental Pratincole | <i>Glareola maldivarum</i> | LC, (1) | | | | | | | | | | | | | | | | | | |
| Little Ringed Plover | <i>Charadrius dubius</i> | LC, (1) | 0.0 | 0.0 | 0.4 | 1.0 | 0.2 | 0.0 | | | | | | V | 0.4 | | V | 0.5 | 0.8 | |
| Kentish Plover | <i>Charadrius alexandrinus</i> | RC, (1) | | | | | | | | | 0.8 | | | | V | | | | | |
| Spotted Redshank | <i>Tringa erythropus</i> | RC, (1) | | | | | | | | | | | | V | | | | | | |
| Common Redshank | <i>Tringa totanus</i> | RC, (1) | | | | | | | | | | | | | | | | | | |
| Marsh Sandpiper | <i>Tringa stagnatilis</i> | RC, (1) | | | | | | | | | | | | | | | | | | |
| Common Greenshank | <i>Tringa nebularia</i> | RC, (1) | V | V | V | | V | | | | | | | | | V | V | | V | |
| Green Sandpiper | <i>Tringa ochropus</i> | (1) | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | | | | | | 1.0 | 0.6 | V | | | 0.5 | |
| Wood Sandpiper | <i>Tringa glareola</i> | LC, (1) | | V | | V | V | | | | | | | V | V | V | V | V | V | |

| Common name | Scientific name | Conservation status | Nov-12 | Dec-12 | Jan-13 | Feb-13 | Mar-13 | Apr-13 | May-13 | Jun-13 | Jul-13 | Aug-13 | Sep-13 | Oct-13 | Nov-13 | Dec-13 | Jan-14 | Feb-14 | Mar-14 | Apr-14 |
|------------------------------|------------------------------------|---------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Common Sandpiper | <i>Actitis hypoleucos</i> | (1) | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.2 | | V | 0.6 | V | 0.8 | 1.6 | 0.2 | V | 0.3 | 1.3 | 0.2 |
| Eurasian Woodcock | <i>Scolopax rusticola</i> | (1) | | | | | | | | | | | | | | | | | | |
| Pintail/Swinhoe's Snipe* | <i>Gallinago stenura/G. megala</i> | LC*,(1) | | | | | | | | | | | | | V | | | | | |
| Common Snipe | <i>Gallinago gallinago</i> | (1) | | V | | | | V | | | | | | | | | | | | |
| Red-necked Stint | <i>Calidris ruficollis</i> | LC, (1) | | | | | | | | | | | | | | | | | | |
| Temminck's Stint | <i>Calidris temminckii</i> | LC, (1) | | | | | | | | | | | | | | | | | | |
| Long-toed Stint | <i>Calidris subminuta</i> | LC, (1) | | | | | | | | | | | | | | | | | | |
| Whiskered Tern | <i>Chlidonias hybrida</i> | (1) | | | | | | | | | | | | | | | | | | |
| Domestic Pigeon | <i>Columba livia</i> | - | | | | | | | | | | | | | | | | | | |
| Oriental Turtle Dove | <i>Streptopelia orientalis</i> | - | | | | | | | | | | | | | | | | | | |
| Red Turtle Dove | <i>Streptopelia tranquebarica</i> | - | | | | | | | | | | | | | | | | | | |
| Spotted Dove | <i>Spilopelia chinensis</i> | - | 0.4 | 1.2 | 0.4 | | 0.2 | 0.2 | 0.4 | 0.5 | 0.8 | 2.8 | 2.0 | 2.6 | 2.6 | 3.0 | 6.2 | 2.3 | 2.0 | 1.0 |
| Asian Koel | <i>Eudynamys scolopaceus</i> | - | | | | | | | | | | | | | | | | | | V |
| Greater Coucal# | <i>Centropus sinensis</i> | | | | | | | | 0.2 | 0.3 | 0.2 | 0.2 | | 0.2 | 0.2 | | | | | 0.4 |
| Eurasian Eagle Owl | <i>Bubo bubo</i> | RC | | | | | | | | | | | V | | V | | | V | | |
| Savanna Nightjar | <i>Caprimulgus affinis</i> | - | | | | V | | | | V | | | | | | | | | V | |
| Pacific Swift | <i>Apus pacificus</i> | (LC) | | | | | | | | | | | | | | | | | | |
| House Swift | <i>Apus nipalensis</i> | - | | 0.4 | V | | | | | | | | | | | | V | 0.5 | | |
| Pied Kingfisher | <i>Ceryle rudis</i> | (LC), (1) | | | | | | V | | | | V | | | | 0.4 | V | 0.3 | V | |
| Common Kingfisher | <i>Alcedo atthis</i> | (1) | 0.2 | 0.2 | 0.5 | 0.4 | 0.2 | 0.2 | 0.6 | 0.3 | 0.4 | 0.4 | 0.8 | 1.0 | 2.0 | 1.6 | 1.2 | 2.3 | 0.8 | 1.0 |
| White-throated Kingfisher | <i>Halcyon smyrnensis</i> | (LC) | | | | | | | | | | | | | V | 0.2 | | | | |
| Barn Swallow | <i>Hirundo rustica</i> | - | 0.2 | 1.0 | | 0.2 | 9.2 | 23.0 | 11.2 | | 1.4 | 1.8 | 1.0 | | V | | | 0.5 | 7.8 | 15.4 |
| Red-rumped Swallow | <i>Hirundo daurica</i> | - | 1.2 | 1.2 | | | | | | | | | | | | | 0.6 | | | |
| Eastern Yellow Wagtail | <i>Motacilla tschutschensis</i> | (1) | 0.2 | 0.2 | 1.4 | 2.8 | 2.2 | 2.2 | 0.2 | | | | | 0.4 | 0.2 | 0.6 | 0.6 | 1.0 | 0.3 | 0.2 |
| Grey Wagtail | <i>Motacilla cinerea</i> | (1) | 0.6 | 0.0 | 0.0 | 0.0 | 0.0 | | | | | | | | | | | | | |
| White Wagtail | <i>Motacilla alba</i> | (1) | 2.6 | 1.0 | 1.0 | 3.0 | 4.8 | 2.0 | 1.0 | | 1.2 | 0.4 | 3.5 | 2.8 | 3.6 | 2.6 | 1.2 | 1.3 | 2.5 | 1.6 |
| Richard's Pipit | <i>Anthus richardi</i> | (1) | 0.2 | 0.4 | 1.2 | 0.4 | 0.4 | | | | | | | | | 0.4 | 0.8 | 2.0 | 2.3 | 3.0 |
| Olive-backed Pipit | <i>Anthus hodgsoni</i> | - | | | | 0.6 | 1.2 | 1.4 | | | | | | | | | | | | 0.2 |
| Red-throated Pipit | <i>Anthus cervinus</i> | LC | | | | | | | | | | | | | | | | | | |
| Buff-bellied Pipit | <i>Anthus rubescens</i> | LC | | | | | | | | | | | | | | | | | | |
| Red-whiskered Bulbul | <i>Pycnonotus jocosus</i> | - | | | | | | 0.2 | | 0.3 | | 0.2 | | | V | | 0.8 | 1.3 | 0.3 | 0.2 |
| Chinese Bulbul | <i>Pycnonotus sinensis</i> | - | | 0.2 | 2.0 | 0.2 | 0.2 | 3.2 | 2.0 | 3.8 | | 2.4 | 3.5 | 0.2 | 1.8 | 4.0 | 3.8 | 0.5 | 0.8 | 1.6 |
| Long-tailed Shrike | <i>Lanius schach</i> | - | 0.2 | | 0.4 | | | 0.2 | | V | 0.4 | 0.2 | 0.3 | 0.2 | | 0.2 | V | 0.3 | | |
| Oriental Magpie Robin | <i>Copsychus saularis</i> | - | | 0.2 | | | | | 0.2 | 0.5 | | 0.2 | 0.3 | V | 0.2 | 0.2 | 0.4 | | | 0.2 |
| Daurian Redstart | <i>Phoenicurus aureoreus</i> | - | | | 0.2 | | | | | | | | | | 0.2 | 0.2 | 0.4 | | | |
| Stejneger's Stonechat | <i>Saxicola stejnegeri</i> | - | 0.4 | 0.4 | 0.8 | | | 0.6 | | | | | | V | 0.2 | 0.2 | 0.4 | 0.5 | 0.3 | V |
| Masked Laughingthrush | <i>Garrulax perspicillatus</i> | - | | | | | 0.6 | 1.2 | | | | | 1.0 | | | 0.4 | 0.8 | V | 0.5 | 0.2 |
| Lanceolated Warbler | <i>Locustella lanceolata</i> | (1) | | | | | | | | | | | | | | | | | | |
| Pallas's Grasshopper Warbler | <i>Locustella certhiola</i> | LC, (1) | | | | | | | | | | | | | | | | | | |
| Black-browed Reed Warbler | <i>Acrocephalus bistrigiceps</i> | (1) | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | | | | | | | | | | | | |
| Zitting Cisticola | <i>Cisticola juncidis</i> | LC | | 0.2 | | | | V | | | | | | | | | 0.4 | | 0.3 | 0.2 |
| Yellow-bellied Prinia | <i>Prinia flaviventris</i> | - | | 0.8 | 0.6 | 0.8 | 1.4 | 4.6 | 4.4 | 5.0 | 2.4 | 1.8 | 3.0 | 2.4 | 0.8 | 1.0 | 0.8 | 1.3 | 2.8 | 0.8 |
| Plain Prinia | <i>Prinia inornata</i> | - | 1.2 | 0.4 | V | 1.4 | 0.2 | 0.2 | 0.2 | 0.5 | 0.8 | 0.4 | 0.3 | | 0.4 | | 0.2 | V | 0.8 | 2.0 |
| Dusky Warbler | <i>Phylloscopus fuscatus</i> | - | | | | | | | | | | | | V | 0.4 | | 0.4 | 0.8 | 1.5 | V |
| Yellow-browed warbler | <i>Phylloscopus inornatus</i> | - | 0.8 | 0.6 | | | | | | | | | | | | | | | | |
| Chinese Penduline-Tit | <i>Remiz consobrinus</i> | RC, (1) | | | | | | | | | | | | | | V | | 4.5 | V | |

| Common name | Scientific name | Conservation status | Nov-12 | Dec-12 | Jan-13 | Feb-13 | Mar-13 | Apr-13 | May-13 | Jun-13 | Jul-13 | Aug-13 | Sep-13 | Oct-13 | Nov-13 | Dec-13 | Jan-14 | Feb-14 | Mar-14 | Apr-14 |
|--------------------------------|----------------------------------|---------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Japanese White-eye | <i>Zosterops japonica</i> | - | | | | | | | | | | | | | | | 0.2 | | | 0.6 |
| Little Bunting | <i>Emberiza pusilla</i> | - | | | | | | | | | | | | | | | | | | |
| Black-faced Bunting | <i>Emberiza spodocephala</i> | - | | | | | | | | | | | | | | | | | V | |
| Chinese Grosbeak | <i>Eophona migratoria</i> | LC | | | | | | | | | | | | | | | | | | |
| Scaly-breasted Munia | <i>Lonchura punctulata</i> | - | | | | 1.0 | 2.2 | 1.0 | | | 2.0 | 0.4 | 0.3 | | | 1.8 | 1.8 | V | 1.8 | V |
| Eurasian Tree Sparrow | <i>Passer montanus</i> | - | | | | | | | | | | | | 0.8 | V | | | | | |
| Red-billed Starling | <i>Spodiopsar sericeus</i> | GC | | | | | | | | | | | | | | | | | V | |
| White-cheeked Starling | <i>Spodiopsar cineraceus</i> | PRC | | | | | | | | | | | | | | | | | | |
| Black-collared Starling | <i>Gracupica nigricollis</i> | - | | | 0.8 | 0.4 | 0.2 | 0.2 | 0.2 | | | | 0.8 | | 0.8 | 1.6 | 0.8 | 0.5 | 1.8 | 0.2 |
| White-shouldered Starling | <i>Sturnia sinensis</i> | (LC) | | | | | | | | | 0.2 | | | | | | | | | |
| Common Myna | <i>Acridotheres tristis</i> | (1) | | | | | | | | | | | | | | | | | | |
| Crested Myna | <i>Acridotheres cristatellus</i> | - | 0.6 | | | | | | 0.2 | 0.3 | 0.8 | 0.2 | | | 2.0 | 1.8 | 1.4 | 0.5 | 1.3 | 1.0 |
| Black-naped Oriole | <i>Oriolus chinensis</i> | LC | | | | | | | | | | | | | | | | | | |
| Black Drongo | <i>Dicrurus macrocercus</i> | - | | | | | | | | | | | 0.3 | 0.4 | V | | | | | V |
| Azure-winged Magpie | <i>Cyanopica cyanus</i> | - | | | | | | | | | | | | | V | | | | | |
| Eurasian Magpie | <i>Pica pica</i> | - | | | | | | | | 0.3 | | | | | V | | 0.8 | V | 0.3 | 0.2 |
| Large-billed Crow | <i>Corvus macrorhynchos</i> | - | | | | | | | | 1.0 | | | | | | | | | | |
| Collared Crow | <i>Corvus torquatus</i> | LC, NT | | | | | | V | | | | | | 0.2 | 0.2 | | | V | 0.5 | 0.8 |
| No. of species recorded | 108 | | | | | | | | | | | | | | | | | | | |

Note: Conservation Status follows that of Fellowes et. al. (2002)

(1) Indicates the bird species is wetland dependent.

* Pintail Snipe and Swinhoe's Snipe cannot be distinguished in field, conservation status refers to Swinhoe's Snipe.

V Indicates the species is recorded outside regular surveys.

DS Indicates species recorded during survey.

Greater Coucal is listed as vulnerable (VU) in China Red Data Book and it is protected under terrestrial wildlife state protection (category II).