

I. Summary of Bird Surveys conducted

Table I1: Summary of bird species of conservation importance and/or wetland-dependence recorded in the Survey Area (excluding the WRA)

Species Name ⁽⁴⁾	Scientific Name ⁽⁴⁾	Wetland Dependence	Conservation Status ⁽¹⁾	Apr 2018 Occurrence ⁽²⁾	Apr 2018 Mean ⁽³⁾	Records outside surveys
Little Grebe	<i>Tachybaptus ruficollis</i>	Y	LC	4	7.0	✓
Grey Heron	<i>Ardea cinerea</i>	Y	PRC	2	0.5	-
Great Egret	<i>Ardea alba</i>	Y	PRC, (RC)	2	0.8	✓
Intermediate Egret	<i>Egretta intermedia</i>	Y	RC	1	-	✓
Little Egret	<i>Egretta garzetta</i>	Y	PRC, (RC)	4	13.8	✓
Eastern Cattle Egret	<i>Bubulcus coromandus</i>	Y	(LC)	1	1.8	✓
Chinese Pond Heron	<i>Ardeola bacchus</i>	Y	PRC, (RC)	4	3.8	✓
Yellow Bittern	<i>Ixobrychus sinensis</i>	Y	(LC)	1	0.3	-
Black-crowned Night Heron	<i>Nycticorax nycticorax</i>	Y	(LC)	1	-	✓
Black Kite	<i>Milvus migrans</i>	Y	(RC)	1	0.3	✓
White-breasted Waterhen	<i>Amaurornis phoenicurus</i>	Y	-	1	0.3	✓
Green Sandpiper	<i>Tringa ochropus</i>	Y	-	1	0.3	✓
Common Sandpiper	<i>Actitis hypoleucos</i>	Y	-	3	3.5	✓
Pied Kingfisher	<i>Ceryle rudis</i>	Y	(LC)	1	0.3	-
Common Kingfisher	<i>Alcedo atthis</i>	Y	-	3	2.0	✓
White Wagtail	<i>Motacilla alba</i>	Y	-	2	1.8	✓
Red-billed Starling	<i>Spodiopsar sericeus</i>	Y	(RC)*	1	0.3	-
White-shouldered Starling	<i>Sturnia sinensis</i>	Y	(LC)	1	-	✓
No. of species recorded:				18		

Note:

(1) Conservation status follows that of Fellowes et al. (2002) and BirdLife International listing (2017) Letters in parentheses indicate that the assessment is on the basis of restrictedness in breeding and/or roosting sites rather than in general occurrence. (Fellowes et al. 2002)

(2) Indicates number of surveys recorded within the reporting period.

(3) Refers to the mean number of individuals recorded in the reporting period (excluding the WRA).

(4) Follows HK bird list (dated 2017-09-05).

* Red-billed Starling is considered by Fellowes et al (2002) to be of Global Concern. Since publication, however, the global population estimate has been revised and the species is not now considered globally threatened. A listing of Regional Concern (RC) based on the importance of the large roosts present near Deep Bay, is considered to be more appropriate. (Wetland Restoration Plan, Mott, 2008). Red-billed Starling is now listed as Least Concern by IUCN. (IUCN, 2016)

Table I2: Summary of bird species of conservation importance and/or wetland-dependence recorded in the WRA

Species Name ⁽⁴⁾	Scientific Name ⁽⁴⁾	Wetland Dependence	Conservation Status ⁽¹⁾	Apr 2018 Occurrence ⁽²⁾	Apr 2018 Mean ⁽³⁾	Records outside surveys
Little Grebe	<i>Tachybaptus ruficollis</i>	Y	LC	3	2.8	✓
Great Cormorant	<i>Phalacrocorax carbo</i>	Y	PRC	-	-	✓
Great Egret	<i>Ardea alba</i>	Y	PRC, (RC)	1	0.3	-
Little Egret	<i>Egretta garzetta</i>	Y	PRC, (RC)	4	4.0	✓
Chinese Pond Heron	<i>Ardeola bacchus</i>	Y	PRC, (RC)	3	1.3	✓
Black Kite	<i>Milvus migrans</i>	Y	(RC)	2	0.5	✓
Eurasian Hobby	<i>Falco subbuteo</i>	Y	(LC)	-	-	✓
White-breasted Waterhen	<i>Amaurornis phoenicurus</i>	Y	-	1	0.3	✓
Common Moorhen	<i>Gallinula chloropus</i>	Y	-	-	-	✓
Common Greenshank	<i>Tringa nebularia</i>	Y	RC	-	-	✓
Green Sandpiper	<i>Tringa ochropus</i>	Y	-	4	1.8	✓
Wood Sandpiper	<i>Tringa glareola</i>	Y	LC	-	-	✓
Common Sandpiper	<i>Actitis hypoleucos</i>	Y	-	2	1.3	✓
Common Kingfisher	<i>Alcedo atthis</i>	Y	-	3	1.0	-
White Wagtail	<i>Motacilla alba</i>	Y	-	1	0.3	✓
White-shouldered Starling	<i>Sturnia sinensis</i>	Y	(LC)	-	-	✓
Collared Crow	<i>Corvus torquatus</i>	Y	LC, NT	-	-	✓
No. of species recorded:				17		

Note:

(1) Conservation status follows that of Fellowes *et al.* (2002) and BirdLife International listing (2017). Letters in parentheses indicate that the assessment is on the basis of restrictedness in breeding and/or roosting sites rather than in general occurrence. (Fellowes *et al.* 2002)

(2) Indicates number of surveys recorded within the reporting period.

(3) Refers to the mean number of individuals recorded in the reporting period in the WRA.

(4) Follows HK bird list (dated 2017-09-05).

* Red-billed Starling is considered by Fellowes *et al.* (2002) to be of Global Concern. Since publication, however, the global population estimate has been revised and the species is not now considered globally threatened. A listing of Regional Concern (RC) based on the importance of the large roosts present near Deep Bay, is considered to be more appropriate. (Wetland Restoration Plan, Mott, 2008). Red-billed Starling is now listed as Least Concern by IUCN. (IUCN, 2016)

Table I3: Summary of bird species recorded in the Survey Area (excluding the WRA) during the reporting month

Species Name ⁽²⁾	Scientific Name ⁽²⁾	Wetland Dependence	Conservation Status ⁽¹⁾
Little Grebe	<i>Tachybaptus ruficollis</i>	Y	LC
Grey Heron	<i>Ardea cinerea</i>	Y	PRC
Great Egret	<i>Ardea alba</i>	Y	PRC, (RC)
Intermediate Egret	<i>Egretta intermedia</i>	Y	RC
Little Egret	<i>Egretta garzetta</i>	Y	PRC, (RC)
Eastern Cattle Egret	<i>Bubulcus coromandus</i>	Y	(LC)
Chinese Pond Heron	<i>Ardeola bacchus</i>	Y	PRC, (RC)
Yellow Bittern	<i>Ixobrychus sinensis</i>	Y	(LC)
Black-crowned Night Heron	<i>Nycticorax nycticorax</i>	Y	(LC)
Black Kite	<i>Milvus migrans</i>	Y	(RC)
White-breasted Waterhen	<i>Amaurornis phoenicurus</i>	Y	-
Green Sandpiper	<i>Tringa ochropus</i>	Y	-
Common Sandpiper	<i>Actitis hypoleucos</i>	Y	-
Spotted Dove	<i>Spilopelia chinensis</i>	N	-
Eurasian Collared Dove	<i>Streptopelia decaocto</i>	N	-
Asian Koel	<i>Eudynamis scolopaceus</i>	N	-
Greater Coucal#	<i>Centropus sinensis</i>	N	-
Pied Kingfisher	<i>Ceryle rudis</i>	Y	(LC)
Common Kingfisher	<i>Alcedo atthis</i>	Y	-
Blue-tailed Bee-eater	<i>Merops philippinus</i>	N	-
Barn Swallow	<i>Hirundo rustica</i>	N	-
White Wagtail	<i>Motacilla alba</i>	Y	-
Olive-backed Pipit	<i>Anthus hodgsoni</i>	N	-
Red-whiskered Bulbul	<i>Pycnonotus jocosus</i>	N	-
Chinese Bulbul	<i>Pycnonotus sinensis</i>	N	-
Long-tailed Shrike	<i>Lanius schach</i>	N	-
Oriental Magpie Robin	<i>Copsychus saularis</i>	N	-
Daurian Redstart	<i>Phoenicurus aureus</i>	N	-
Masked Laughingthrush	<i>Garrulax perspicillatus</i>	N	-
Common Tailorbird	<i>Orthotomus sutorius</i>	N	-
Scaly-breasted Munia	<i>Lonchura punctulata</i>	N	-
Eurasian Tree Sparrow	<i>Passer montanus</i>	N	-
Red-billed Starling	<i>Spodiopsar sericeus</i>	Y	(RC)*
Black-collared Starling	<i>Gracupica nigricollis</i>	N	-
White-shouldered Starling	<i>Sturnia sinensis</i>	Y	(LC)
Crested Myna	<i>Acridotheres cristatellus</i>	N	-
Eurasian Magpie	<i>Pica pica</i>	N	-
No. of species recorded:		37	

Note:

- (1) Conservation status follows that of Fellowes *et al.* (2002) and BirdLife International listing (2017). Letters in parentheses indicate that the assessment is on the basis of restrictedness in breeding and/or roosting sites rather than in general occurrence. (Fellowes *et al.* 2002)
- (2) Follows HK bird list (dated 2017-09-05)
- # Greater Coucal is listed as vulnerable (VU) in China Red Data Book and it is protected under terrestrial wildlife state protection (category II)
- * Red-billed Starling is considered by Fellowes *et al.* (2002) to be of Global Concern. Since publication, however, the global population estimate has been revised and the species is not now considered globally threatened. A listing of Regional Concern (RC) based on the importance of the large roosts present near Deep Bay, is considered to be

more appropriate. (Wetland Restoration Plan, Mott, 2008). Red-billed Starling is now listed as Least Concern by IUCN. (IUCN, 2016)

Table I4: Summary of bird species recorded in the WRA during the reporting month

Species Name ⁽²⁾	Scientific Name ⁽²⁾	Wetland Dependence	Conservation Status ⁽¹⁾
Little Grebe	<i>Tachybaptus ruficollis</i>	Y	LC
Great Cormorant	<i>Phalacrocorax carbo</i>	Y	PRC
Great Egret	<i>Ardea alba</i>	Y	PRC, (RC)
Little Egret	<i>Egretta garzetta</i>	Y	PRC, (RC)
Chinese Pond Heron	<i>Ardeola bacchus</i>	Y	PRC, (RC)
Black Kite	<i>Milvus migrans</i>	Y	(RC)
Eurasian Hobby	<i>Falco subbuteo</i>	Y	(LC)
White-breasted Waterhen	<i>Amaurornis phoenicurus</i>	Y	-
Common Moorhen	<i>Gallinula chloropus</i>	Y	-
Common Greenshank	<i>Tringa nebularia</i>	Y	RC
Green Sandpiper	<i>Tringa ochropus</i>	Y	-
Wood Sandpiper	<i>Tringa glareola</i>	Y	LC
Common Sandpiper	<i>Actitis hypoleucos</i>	Y	-
Spotted Dove	<i>Spilopelia chinensis</i>	N	-
Eurasian Collared Dove	<i>Streptopelia decaocto</i>	N	-
Asian Koel	<i>Eudynamis scolopaceus</i>	N	-
Greater Coucal#	<i>Centropus sinensis</i>	N	-
Common Kingfisher	<i>Alcedo atthis</i>	Y	-
Barn Swallow	<i>Hirundo rustica</i>	N	-
White Wagtail	<i>Motacilla alba</i>	Y	-
Olive-backed Pipit	<i>Anthus hodgsoni</i>	N	-
Red-whiskered Bulbul	<i>Pycnonotus jocosus</i>	N	-
Chinese Bulbul	<i>Pycnonotus sinensis</i>	N	-
Long-tailed Shrike	<i>Lanius schach</i>	N	-
Oriental Magpie Robin	<i>Copsychus saularis</i>	N	-
Masked Laughingthrush	<i>Garrulax perspicillatus</i>	N	-
Common Tailorbird	<i>Orthotomus sutorius</i>	N	-
Dusky Warbler	<i>Phylloscopus fuscatus</i>	N	-
Yellow-bellied Prinia	<i>Prinia flaviventris</i>	N	-
Scaly-breasted Munia	<i>Lonchura punctulata</i>	N	-
Eurasian Tree Sparrow	<i>Passer montanus</i>	N	-
Black-collared Starling	<i>Gracupica nigricollis</i>	N	-
White-shouldered Starling	<i>Sturnia sinensis</i>	Y	(LC)
Crested Myna	<i>Acridotheres cristatellus</i>	N	-
Eurasian Magpie	<i>Pica pica</i>	N	-
Collared Crow	<i>Corvus torquatus</i>	Y	LC, NT
No. of species recorded:		36	

Note:

- (1) Conservation status follows that of Fellowes *et al.* (2002) and BirdLife International listing (2017). Letters in parentheses indicate that the assessment is on the basis of restrictedness in breeding and/or roosting sites rather than in general occurrence. (Fellowes *et al.* 2002)
- (2) Follows HK bird list (dated 2017-09-05)
- # Greater Coucal is listed as vulnerable (VU) in China Red Data Book and it is protected under terrestrial wildlife state protection (category II)

- * Red-billed Starling is considered by Fellows et al (2002) to be of Global Concern. Since publication, however, the global population estimate has been revised and the species is not now considered globally threatened. A listing of Regional Concern (RC) based on the importance of the large roosts present near Deep Bay, is considered to be more appropriate. (Wetland Restoration Plan, Mott, 2008). Red-billed Starling is now listed as Least Concern by IUCN. (IUCN, 2016)